

UNITED STATES
SECURITIES AND EXCHANGE COMMISSION
Washington, D.C. 20549

FORM 8-K

CURRENT REPORT
Pursuant to Section 13 or 15(d) of
the Securities Exchange Act of 1934

Date of Report (Date of earliest event reported): June 24, 2024

AVALO THERAPEUTICS, INC.
(Exact name of registrant as specified in its charter)

Delaware

(State or other jurisdiction of incorporation)

001-37590
(Commission File Number)

45-0705648
(IRS Employer Identification No.)

540 Gaither Road, Suite 400, Rockville, Maryland 20850
(Address of principal executive offices) (Zip Code)

Registrant's Telephone Number, Including Area Code: (410) 522-8707

Check the appropriate box below if the Form 8-K filing is intended to simultaneously satisfy the filing obligation of the registrant under any of the following provisions:

- ☐ Written communications pursuant to Rule 425 under the Securities Act (17 CFR 230.425)
- ☐ Soliciting material pursuant to Rule 14a-12 under the Exchange Act (17 CFR 240.14a-12)
- ☐ Pre-commencement communications pursuant to Rule 14d-2(b) under the Exchange Act (17 CFR 240.14d-2(b))
- ☐ Pre-commencement communications pursuant to Rule 13e-4(c) under the Exchange Act (17 CFR 240.13e-4(c))

Securities registered pursuant to Section 12(b) of the Act:

Title of each class	Trading Symbol(s)	Name of each exchange on which registered
Common Stock, \$0.001 Par Value	AVTX	Nasdaq Capital Market

Indicate by check mark whether the registrant is an emerging growth company as defined in Rule 405 of the Securities Act of 1933 (§230.405 of this chapter) or Rule 12b-2 of the Securities Exchange Act of 1934 (§240.12b-2 of this chapter).

Emerging Growth Company ☐

If an emerging growth company, indicate by check mark if the registrant has elected not to use the extended transition period for complying with any new or revised financial accounting standards provided pursuant to Section 13(a) of the Exchange Act. ☐

Item 8.01 Other Events.

On June 24, 2024, Avalo Therapeutics, Inc. (the “Company”) posted on its website an updated investor presentation (the “Investor Presentation”). The Investor Presentation will be used from time to time in meetings with investors. A copy of the Investor Presentation is attached hereto as Exhibit 99.1 and is incorporated herein by reference.

Item 9.01 Financial Statements and Exhibits.

(d) Exhibits:

Exhibit No.	Description
99.1	Investor Presentation.
104	The cover pages of this Current Report on Form 8-K, formatted in Inline XBRL.

SIGNATURE

Pursuant to the requirements of the Securities Exchange Act of 1934, the registrant has duly caused this report to be signed on its behalf by the undersigned hereunto duly authorized.

AVALO THERAPEUTICS, INC.

Date: June 24, 2024

By: /s/ Christopher Sullivan

Christopher Sullivan
Chief Financial Officer

Avalo Therapeutics, Inc. (AVTX)

Corporate Presentation

June 2024




avalo
THERAPEUTICS

Forward-Looking Statements

This presentation may include forward-looking statements made pursuant to the Private Securities Litigation Reform Act of 1995. Forward-looking statements are statements that are not historical facts. Such forward-looking statements are subject to significant risks and uncertainties that are subject to change based on various factors (many of which are beyond Avalo's control), which could cause actual results to differ from the forward-looking statements. Such statements may include, without limitation, statements with respect to Avalo's plans, objectives, projections, expectations and intentions and other statements identified by words such as "projects," "may," "might," "will," "could," "would," "should," "continue," "seeks," "aims," "predicts," "believes," "expects," "anticipates," "estimates," "intends," "plans," "potential," or similar expressions (including their use in the negative), or by discussions of future matters such as: integration of AVTX-009 into our operations; drug development costs, timing of trial results and other risks, including reliance on investigators and enrollment of patients in clinical trials; the intended use of the proceeds from the private placement; reliance on key personnel; regulatory risks; general economic and market risks and uncertainties, including those caused the war in Ukraine and the Middle East; and those other risks detailed in Avalo's filings with the Securities and Exchange Commission, available at www.sec.gov. Actual results may differ from those set forth in the forward-looking statements. Except as required by applicable law, Avalo expressly disclaims any obligations or undertaking to release publicly any updates or revisions to any forward-looking statements contained herein to reflect any change in Avalo's expectations with respect thereto or any change in events, conditions or circumstances on which any statement is based.

Avalo Therapeutics & AlmataBio: A Strong Strategic Fit to Drive Long-Term Growth



Concurrent with the acquisition of AVTX-009 in March 2024, Avalo announced a PIPE financing of up to \$185M providing cash runway into 2027

Avalo Therapeutics Pipeline

Compound	Indication	PreClin	P1	P2	P3	Anticipated Milestones
AVTX-009 Anti-IL-1 β mAb	Hidradenitis suppurativa (HS)					P2 Topline Results 2026
	Autoimmune Indication TBD					TBD
Next Generation IL-1 β (extended half-life)	--					TBD
Quisovalimab AVTX-002, Anti-LIGHT mAb	--					<i>Under strategic review</i>
AVTX-008 BTLA agonist fusion protein	--					<i>Under strategic review</i>

Executive Summary and Development Timeline

- **Potential for a Best-in-Disease Profile in HS**

- High potency and favorable half-life may allow for improved efficacy and convenient dosing
- Potential in other autoimmune diseases

- **Key Clinical Evidence Supporting IL-1 β in HS**

- In a large, well controlled Phase 2 trial (NCT05139602), lutikizumab validates IL-1 β targeting in HS. Efficacy was comparable with other HS therapies despite a more severe patient population¹
- Clinical evidence suggests anti-IL-1 α therapy is not effective in HS^{2,3}
- MAS825 (IL-1 β /IL-18 bispecific) showed positive results in a Phase 2 randomized controlled study (NCT03827798)⁴
- Monospecific IL-1 β inhibition may outperform bispecifics that address targets that are unvalidated (IL-18) or known not to contribute to efficacy (IL-1 α)
- We believe AVTX-009 has a high probability of success in HS

- **HS Anticipated to Become Multi-Billion Dollar Market**

- **HS Topline Results Expected in 2026**

- **Expected Cash Runway into 2027**

1. Kimball AB, et al. Presented at: American Academy of Dermatology; March 8-12, 2024; San Diego, CA
2. ClinicalTrials.gov identifier: NCT04988308. Updated November 13, 2023. Accessed March 24, 2024. <https://clinicaltrials.gov/search?term=NCT04988308> NCT04019041
3. ClinicalTrials.gov identifier: NCT04019041. Updated July 27, 2023. Accessed March 24, 2024. <https://clinicaltrials.gov/search?term=NCT04988308>
4. Kimball AB, et al. Presented at: American Academy of Dermatology; March 8-12, 2024; San Diego, CA

Experienced Management Team

Decades of successful leadership, product development, and commercialization in pharma and biotech



Garry A. Neil, MD
Chief Executive Officer
Chairman of the Board



Chris Sullivan
Chief Financial Officer



Lisa Hegg, PhD
SVP, Program Management,
Corporate Infrastructure,
Clinical Operations



Colleen Matkowski
SVP, Global Regulatory Affairs,
Quality Assurance



Dino C. Miano, PhD
SVP, CMC,
Technical Operations



GlaxoSmithKline




AVTX-009


Highly Potent and Specific Inhibitor of a Validated Immune Target; Potential for Q2W to Q12W Dosing^{1,2}

- **High-affinity humanized antibody that potently neutralizes IL-1 β**
 - Originally developed by Lilly
 - Exceptional K_d of <3 pM (picomolar)
 - Superior potency vs ILARIS in vitro
 - t_{1/2} ~19 days (SC and IV)
 - Bioavailability ~73%
- **Clinical experience: 245 patients studied in Phase 1 & Phase 2 trials³**
 - Excellent tolerability and safety at all doses up to 180 mg weekly
 - Marked lowering of hs-CRP after a single dose
 - Potency and half life expected to support Q4W or less frequent dosing in HS
- **Suitable for subcutaneous and intravenous formulation**
 - Stable 150 mg/ml dosage form
 - Plan for commercial presentation to be an autoinjector

AVTX-009 has Higher Potency than Canakinumab and Gevokizumab



AVTX-009 has Strong Pharmacokinetic Profile




1. Data on file
 2. Bihorel S, et al. *AAPS J* 2014;16(5):1009-1017
 3. Sloan-Lancaster J, et al. *Diabetes Care* 2013;36(8):2239-2246



IL-1 β is a Validated Target in Inflammatory Diseases

- IL-1 β is a central driver of the inflammatory process¹: activates immune cells that generate pro-inflammatory cytokines including IL-6, TNF- α , and IL-17
- Inhibition of IL-1 β has been shown to be effective and safe in a variety of inflammatory diseases including Hidradenitis Suppurativa (HS)²
- IL-1 β is involved in the pathogenesis of many autoimmune and autoinflammatory diseases




1. Dinarello CA. *Immunol Rev*. 2018;281(1):8-27
2. Kany S, et al. *Int J Mol Sci*. 2019;20(23):6008

Hidradenitis Suppurativa (HS)

avalo
THERAPEUTICS

Hidradenitis Suppurativa (HS)

- Chronic, often debilitating inflammatory skin disease
 - Lumps, abscesses and scars develop under the arms, in the groin and other areas
- Current treatments:
 - Antibiotics
 - Retinoids
 - Steroids-topical, oral, injections
 - Cosentyx, Humira
- HS has an estimated prevalence of 0.7–1.2% in the European-US population¹




Hurley Stages 1-3 (a-c) and Scarring folliculitis (d)

1. Nguyen TV, et al. *J Eur Acad Dermatol Venerol*. 2021;35(1):50-61

2. Ngan V, et al. Hidradenitis suppurativa. *DermNet*. Accessed February 16, 2024. <https://dermnetnz.org/topics/hidradenitis-suppurativa>

IL-1 β is Strongly Implicated in the Pathophysiology of HS¹

- Inflammatory cascade in HS is triggered by various external stimuli
 - Smoking, dysbiosis, or mechanical stress
- IL-1 β is a key driver of the inflammatory cascade that leads to the destruction of the pilosebaceous unit
 - Increased IL-1 β levels in lesional skin^{2,3}
 - Genetic associations⁴
 - Clinical benefit has been observed with anti-IL-1 drugs



DAMPs, damage-associated molecular pattern molecules; DC, dendritic cell; IL, interleukin; IL-R, interleukin receptor; PAMPs, pathogen-associated molecular pattern molecules.

1. Calabrese L, et al. *Biomolecules*. 2024;14(2):175
 2. Vossen ARJV, et al. *J Invest Dermatol*. 2020;140(7):1463-1466.e2
 3. Kelly G, et al. *Br J Dermatol*. 2015;173(6):1431-1439
 4. Marzano AV, et al. *Dermatology*. 2022;238(5):860-869

Baseline Patient Characteristics of Biologics from Recent HS Trials

Lutikizumab Trial Enrolled More Severe Patients than Competitor Trials

Patient Characteristics	adalimumab PIONEER I / II ¹	secukinumab SUNSHINE / SUNRISE ²	bimekizumab BE HEARD I / II ³	sonelokimab MIRA ⁴	lutikizumab NCT05139602 ⁵
Age (years), mean	34.9 – 37.8	35.5 – 37.3	36.7 / 36.6	37.6	37.0-39.5
Gender, female, %	59.5 – 69.3	54 – 57	63.0 / 50.7	59.8	53.8-67.6
Race, White, %	75.8 – 87.7	74 – 81	77.8 / 81.5	85.0	64.9-88.9
BMI, kg/m ² , mean	31.3 – 34.5	31.4 – 32.8	33.8 / 32.3	33.7	33.0-34.1
Smoking, current, %	52.9 – 67.3	50 – 58	43.0 / 48.1	46.6	24.3-46.2
Duration of HS years, mean	8.8 – 9.9	6.6 – 8.2	9.0 / 7.0	8.5	10.0-13.2
Lesions, mean					
- AN count	10.7 – 14.4	12.6 – 13.9	16 / 16.5	14.0	11.4-17.0
- DT	3.0 – 4.6	3.2 – 3.6	3.8 / 3.4	3.5	5.7-8.7
Hurley stage, %					
- I	0	2 – 6	0	0	
- II	52.3 – 54.6	51 – 60	50.3 / 61.1	63.7	25.6-35.1
- III	45.4 – 47.7	28 – 46	49.7 / 38.9	36.3	64.9-74.4
Prior biologic use, %	0	20 - 26	25.0 / 13.2	17.5	100 TNF failure entry criteria

1. Kimball AB et al. N Engl J Med. 2016; 375:422-34;

2. Kimball AB et al. Lancet. 2023; 401:747-7613;


3. Presentations at American Academy of Dermatology (AAD 2023), March 17-21 New Orleans, LA and 2023 European Academy of Dermatology and Venereology (EADV 2023), October 11-14, Berlin, Germany

4. R&D Presentation – Results MIRA trial June 26th 2023 <https://ir.moonlakeco.com/static-files/86c71a51-5836-4f1c-9a2c-45e440a50d75>

5. Kimball AB et al. presentation at American Academy of Dermatology (AAD 2024), 8-12 March 2024, San Diego, CA


Lutikizumab Efficacy Comparable to Other Agents in a More Severe Patient Population that Failed TNF- α Therapy¹

HiSCR75



Lower response rates observed in patients that have been previously treated with biologic agents or Hurley Stage 3²⁻⁴

HiSCR50



1. Kimball AB, et al. Presented at: American Academy of Dermatology; March 8-12, 2024; San Diego, CA
2. Kimball AB, et al. Supplement. *N Engl J Med*. 2016;375(5):422-434
3. Sayed, et al. Poster presented at: European Academy of Dermatology and Venereology Congress; October 11-14, 2023. Berlin, Germany
4. Zouboulis CC, et al. *Br J Dermatol*. Published online March 12, 2024. doi:10.1093/bjd/ljae098

IL-1 α is Unlikely to be an Important Driver of HS Pathophysiology

- **Phase 2, bermekimab (IL-1 α mAb) (NCT04019041)¹**
 - Moderate to Severe Hidradenitis Suppurativa
 - Loading doses and then either 400 mg QW or 400 mg Q2W
 - Primary endpoint: HiSCR50 at week; Efficacy not demonstrated
- **Phase 2, bermekimab (NCT04988308)²**
 - Moderate to Severe Hidradenitis Suppurativa
 - Primary endpoint : HiSCR50 at week 16
 - **Study was terminated prematurely as interim analysis met the futility criteria for primary endpoint (no better than placebo)**


	Part 1: Placebo (N=35)	Part 1: Bermekimab (N=35)	Part 1: Adalimumab (N=35)
Primary Endpoint Percentage of Participants Who Achieved Hidradenitis Suppurativa Clinical Response-50 (HiSCR50) at Week 16	37.1	37.1	57.1
Key Secondary Endpoint Percentage of Participants Who Achieved HiSCR75 at Week 16	25.7	25.7	40.0

1. ClinicalTrials.gov identifier: NCT04988308. Updated November 13, 2023. Accessed March 24, 2024. <https://clinicaltrials.gov/search?term=NCT04988308> NCT04019041
2. ClinicalTrials.gov identifier: NCT04019041. Updated July 27, 2023. Accessed March 24, 2024. <https://clinicaltrials.gov/search?term=NCT04988308>

MAS825 May Provide Further Clinical Evidence for IL-1 β in HS

NCT03827798: Phase 2b Platform study in moderate to severe HS

- Novartis' MAS825 is a bispecific inhibitor of IL-1 β and IL-18
- MAS825 arm succeeded in a placebo-controlled platform trial in HS^{1,2}
- To our knowledge, there is no other clinical validation of IL-18 inhibition in HS
- Unlike IL-1 β , IL-18 is not strongly implicated in the pathophysiology of HS



Patients


- Adult patients aged 18–65 years
- Moderate to severe HS for ≥ 12 months in ≥ 2 anatomical areas with ≤ 15 tunnels
- **Cohorts A, C, and E:** ≥ 5 inflammatory lesions
- **Cohorts B and D:** ≥ 3 inflammatory lesions

1. ClinicalTrials.gov identifier: NCT03827798. Updated January 21, 2023. Accessed March 24, 2024. <https://clinicaltrials.gov/search?term=NCT03827798>

2. Kimball AB, et al. Presented at: American Academy of Dermatology; March 8–12, 2024; San Diego, CA

Large Unmet Need in HS

Global HS market has multibillion dollar sales potential with sponsors of several approved products projecting annual sales of \$1.5 – 3B¹




1. Independent Market Analysis by Biotech Value Advisors LLC
2. Nguyen TV, et al. *J Eur Acad Dermatol Venereol.* 2021;35(1):50-61
3. Garg A, et al. *JAMA Dermatol.* 2017;153(8):760-764
4. UpToDate as of 02/15/24

Phase 2 Study in Hidradenitis Suppurativa (AVTX-009-HS-201)

Efficacy and Safety of AVTX-009 Treatment in Participants With Hidradenitis Suppurativa (HS)

Key Inclusion Criteria

- Signs and symptoms of HS for at least 6 months prior to baseline as determined by the investigator.
- Total abscess and inflammatory nodule (AN) count of ≥ 5 at baseline AND HS lesions must be present in at least 2 distinct anatomic areas, at least one of which is Hurley Stage 2 or 3.
- Number of patients who have not failed anti-TNF therapy [anti-TNF naïve or exposed but not failure] will be limited to approximately 40%, the remainder must have failed anti-TNF treatment in the opinion of investigator.



Primary Study Endpoint

Primary Endpoint: Percentage of Participants Achieving Hidradenitis Suppurative Clinical Response HiSCR75 at 16 weeks defined as:

At least a 75% reduction in the total abscess and inflammatory nodule (AN) count, with no increase in abscess count and no increase in draining fistula count relative to Baseline

Key Secondary/Exploratory Endpoints

Key Secondary Endpoints:

- Adverse Events (AEs) and tolerability
- HiSCR50, HiSCR90
- International HS Severity Score System (IHS4)
- AN Count, Draining Fistula Count
- Patient's Global Assessment of Skin Pain (PGA Skin Pain) (NRS30)
- Percentage of subjects with flares
- ADA

Exploratory Endpoints:

- PK
- HISQOL, DLQI, PHQ-9
- Biomarkers- CRP, IL-6, potentially other biomarkers


Trial has 80% power to show a HiSCR75 response for each individual arm (based on lutikizumab Phase 2 HiSCR75)

Potential Additional Indications

avalo
THERAPEUTICS

Role of IL-1 β in IBD

- IL-1 β plays a central role in inflammation in IBD¹
 - IL-1 β being a key cytokine produced upon inflammasome activation
 - Dysregulated inflammasome activation has been implicated in the pathogenesis of Crohn's Disease (CD)
- IL-1-driven stromal–neutrophil interactions define a subset of patients that do not respond to current therapies ^{2,3}
- Observed overlap of patients that have IBD and HS^{4,5}



1. Mao L, et al. *Front Immunol*. 2018;9:2566
 2. Friedrich M, et al. *Nat Med*. 2021;27(11):1970-1981
 3. Cader MZ, Kaser A. *Nat Med*. 2021;27(11):1870-1871
 4. Chen WT, Chi CC. *JAMA Dermatol*. 2019;155(9):1022-1027
 5. Zhang M, et al. *World J Clin Cases*. 2021;9(15):3506-3516

Recent IL-1 Trial Initiations in IBD

- The goal of IBD therapeutics is remission
 - Only a minority of IBD patients obtain remission with current therapies
- AbbVie plans to evaluate lutikizumab, dual-variable-domain interleukin (IL) 1 α /1 β antagonist as monotherapy in UC and in combination with SKYRIZI in Crohn's
 - "...we believe lutikizumab has the potential to be used in combinations to provide transformational levels of efficacy in IBD. We plan to evaluate combo approaches with lutikizumab and Skyrizi... in Crohn's. Our Phase 2 studies in IBD are expected to begin later this year."--Roopal Thakkar, Senior Vice President, Chief Medical Officer, Global Therapeutics --from AbbVie 4Q23 Earnings Call Transcript
- There is an opportunity for greater efficacy for patients with IBD with anti-IL-1 β as a monotherapy and in combination

Executive Summary

The logo for Avalo Therapeutics features a large, teal-colored, hand-drawn circular brushstroke that is open on the right side. The word "avallo" is written in a white, lowercase, sans-serif font, with the "a" and "o" having a slight shadow or double outline. Below "avallo", the word "THERAPEUTICS" is written in a smaller, white, uppercase, sans-serif font.

avallo
THERAPEUTICS

Executive Summary and Development Timeline

- **Potential for a Best-in-Disease Profile in HS**

- High potency and favorable half-life may allow for improved efficacy and convenient dosing
- Potential in other autoimmune diseases

- **Key Clinical Evidence Supporting IL-1 β in HS**

- In a large, well controlled Phase 2 trial (NCT05139602), lutikizumab validates IL-1 β targeting in HS. Efficacy was comparable with other HS therapies despite a more severe patient population¹
- Clinical evidence suggests anti-IL-1 α therapy is not effective in HS^{2,3}
- MAS825 (IL-1 β /IL-18 bispecific) showed positive results in a Phase 2 randomized controlled study (NCT03827798)⁴
- Monospecific IL-1 β inhibition may outperform bispecifics that address targets that are unvalidated (IL-18) or known not to contribute to efficacy (IL-1 α)
- We believe AVTX-009 has a high probability of success in HS

- **HS Anticipated to Become Multi-Billion Dollar Market**

- **HS Topline Results Expected in 2026**

- **Expected Cash Runway into 2027**

1. Kimball AB, et al. Presented at: American Academy of Dermatology; March 8-12, 2024; San Diego, CA
2. ClinicalTrials.gov identifier: NCT04988308. Updated November 13, 2023. Accessed March 24, 2024. <https://clinicaltrials.gov/search?term=NCT04988308> NCT04019041
3. ClinicalTrials.gov identifier: NCT04019041. Updated July 27, 2023. Accessed March 24, 2024. <https://clinicaltrials.gov/search?term=NCT04988308>
4. Kimball AB, et al. Presented at: American Academy of Dermatology; March 8-12, 2024; San Diego, CA

Appendix

The logo for Avalo Therapeutics is located in the bottom right corner of the slide. It features a large, stylized, light blue circular graphic that is partially cut off by the right edge of the slide. The word "avallo" is written in a white, lowercase, sans-serif font, and the word "THERAPEUTICS" is written in a smaller, white, uppercase, sans-serif font directly below it.

avallo
THERAPEUTICS

Financial & Investor Information

NASDAQ: AVTX

The following data is as of March 31, 2024

- Cash and cash equivalents – \$110.2M
- Expected cash runway into 2027
- Outstanding common shares – 1.0M
- Fully diluted shares – 35.4M¹

¹ Based on 1.0 million shares of common stock outstanding, 22.4 million shares of common stock underlying outstanding convertible non-voting Series C preferred stock, 12.0 million shares of common stock outstanding warrants and 7,513 shares of common stock underlying outstanding stock options. Subject to and upon Avalo stockholder approval, each share of convertible non-voting Series C preferred stock will automatically convert into 1,000 shares of Avalo common stock, subject to certain beneficial ownership limitations. Approximately 12.0 million warrants were issued on March 28, 2024 that are exercisable for approximately \$5.80 per underlying share of common stock for an aggregate of up to \$69.4 million of gross proceeds. These warrants are exercisable for common stock, or an equivalent amount (as converted to common stock) of non-voting convertible preferred stock in the event that certain beneficial ownership limitation applicable to each holder is reached, until the earlier of five years from issuance or 30 days after the public announcement of the first patient dosed in a Phase 2 trial of AVTX-009 in hidradenitis suppurativa.

NASDAQ:AVTX

www.avalotx.com

The logo for Avalo Therapeutics features a large, teal-colored, hand-drawn style arc that forms a partial circle on the right side of the slide. The word "avalotx" is written in a white, lowercase, sans-serif font, with the "o" being slightly larger and more rounded. Below it, the word "THERAPEUTICS" is written in a smaller, white, uppercase, sans-serif font.

avalotx
THERAPEUTICS

